



University Orthopedics, Inc.

BICEP TENODESIS REHABILITATION PROTOCOL

This protocol was developed for patients who have had a bicep tenodesis. Please note this protocol is a <u>guideline</u>. Patients with additional surgery will progress at different rates. Achieving the criteria of each phase should be emphasized more than the approximate duration.

PHASE I: ~1-7 DAYS POSTOPERATIVE

GOALS:

- Pain control
- Protection

AMBULATION AND BRACE USE: Place pillow under shoulder / arm while sleeping for comfort NO RESISTED Shoulder Abduction/Flexion or Elbow Flexion/Supination x 6 wks.

EXERCISES:

Hand squeezing exercises Elbow and wrist active motion (AROM) with shoulder in neutral position at side Supported pendulum exercises Shoulder shrugs / scapular retraction without resistance Ice pack

PHASE II: ~1-6 WEEKS POSTOPERATIVE

GOALS:

Full AROM

AMBULATION AND BRACE USE: Continue sling x 6 weeks

EXERCISES: Full pendulum exercises AROM/AAROM as tolerated (pulleys/wand/wall walks) 1-2 Finger Isometrics x 6 (fist in box) Stationary bike (must wear sling)

PHASE III: ~6-9 WEEKS POSTOPERATIVE

GOALS:

• 30 Wall Push-ups

AMBULATION AND BRACE USE: D/C Sling

EXERCISES:

Continue appropriate previous exercises Rotator cuff strengthening with Theraband – ER and IR arm at side with rolled towel in axilla

- Flexion to 60 degrees
- Abduction to 60 degrees
- Scaption to 60 degrees
- Extension to 30 degrees
- Standing rows with Theraband
- Resistive elbow / wrist exercises with light dumbbell
- Push-up progression Wall, etc
- Body Blade
- Ball on wall (arcs, alphabet)
- BAPS / BOSU on hands
- UBE Forwards and backwards at low
- resistance
- Elliptical trainer
- Stairmaster
- Pool walking / running No UE resistive exercises

PHASE IV: ~9-12 WEEKS POSTOPERATIVE GOALS:







Rotator cuff strength WNL

EXERCISES:

Continue appropriate previous exercises with increased resistance

PROM / mobilization as needed to regain full ROM

Seated row weight machine with light weight Push-up progression – Wall to table Ball toss with arm at side using light ball Treadmill – Running progression program Pool therapy – With UE resistance

PHASE V: ~3-4 MONTHS POSTOPERATIVE

GOALS:

 Run 2 miles at easy pace, 30 chair push-ups

EXERCISES:

Continue appropriate previous exercises Fitter on hands Ball toss overhead Weight training with light resistance (no overhead press, pull downs, lateral lifts) Push-up progression – Table to chair

PHASE VI: ~4-6 MONTHS POSTOPERATIVE

GOALS:

• Resume all activities

EXERCISES: Continue appropriate previous exercises Push-ups, regular Sit-ups Gravitron – Pull-ups and dips Swimming Running progression to track Progressive weight training program Transition to home / gym program

No contact sports until 6 months post-op

**This protocol is designed to be administered by a licensed physical therapist and/or certified athletic trainer. Please do not hesitate to contact our office should you have any questions concerning the rehabilitation process.